

PRODUCT DATA SHEET

Volatile Acid Mixture (qualitative)

Catalog No: 1075
Solvent: DI Water
Storage: 4-8°C
Concentration: various
Quantity: 100ml

GC Conditions:

Column: Nukol 30m x 0.53mm
Carrier Gas: helium
Make-up Gas: helium
Split Ratio: 10:1
Oven Initial: 100°C
Oven Final: 200°C
Detector: FID, 230 °C

Linear Velocity: 17cm/sec
Flow Rate: 40ml/min
Vent Flow: 70ml/min
Program Rate: 8°C/min
Hold Time: 2.5 min
Injector: 230°C

Components:

- Formic acid
- Acetic acid
- Propionic acid
- Isobutyric acid
- N-butyric acid
- Isovaleric acid
- N-valeric acid
- Isocaproic acid
- N-caproic acid
- Heptanoic acid

Application Notes:

This mixture contains ten volatile fatty acids and is ideal for their identification by gas chromatography, mass spectrometry, and high performance liquid chromatography and is prepared from high purity stock materials. Knowledge of the fatty acid content of bacteria, for example, can be of great benefit in understanding microbials and can be of great nutritional importance in animals and humans.^{1,2,3} This is a qualitative mixture and should not be used for quantitative purposes.

Selected References:

1. M. Or-Rashid, N. Odongo and B. McBride, "Fatty acid composition of ruminal bacteria and protozoa, with emphasis on conjugated linoleic acid, vaccenic acid, and odd-chain and branched-chain fatty acids" *Journal of Animal Science*, Vol. 85 pp. 1228, 2007
2. Y. Zhang, S. White, and C. Rock "Inhibiting Bacterial Fatty Acid Synthesis" *The Journal of Biological Chemistry*, Vol. 281(26) pp. 17541, 2006
3. N. Rozès et al. "A rapid method for the determination of bacterial fatty acid composition" *Applied Microbiology*, Vol. 3(17) pp. 126, 1993

This product is to be used for research only. It is not intended for drug or diagnostic use, human consumption or to be used in food or food additives. Matreya assumes no liability for any use of this product by the end user. We believe the information, offered in good faith, is accurate.