PRODUCT DATA SHEET



N-Hexadecanoyl-phytosphingosine

Catalog number: 2035

Common names: N-C16:0-Phytoceramide; N-

Palmitoyl-phytosphingosine

Source: semisynthetic, yeast (*Pichia ciferri*)

Solubility: chloroform/methanol, 5:1

CAS number: 11149-09-8

Molecular Formula: C₃₄H₆₉NO₄

Molecular Weight: 556

Storage: -20°C

Purity: TLC >98%, HPLC >98%; identity

confirmed by MS

TLC System: chloroform/methanol (90:10 by vol.)

Appearance: solid

Application Notes:

Phytosphingosine is a long-chain sphingoid base having important cellular functions such as signaling, cytoskeletal structure, cellular cycle, and heat stress response. It is found largely in mammals, plants, and yeast. Phytosphingosine has seen much use in cosmetics due to its effects on the skin such as reducing inflammation by inhibiting the expression of the allergic cytokines IL-4 and TNF-α and the activation of the transcription factors NF-jB and c-jun in histamine-stimulated skin tissues.¹ Phytosphingosine can lead to apoptosis via two distinct pathways and has been investigated as a possible cancer therapeutic treatment. Phytoceramides (fatty acid acylated to Phytosphingosine) are distributed at the microvillous membrane of the epithelial cells of the small intestine. Crypt cells and the adjacent epithelial cells produce phytosphingoglycolipids in much greater quantities than more differentiated epithelial cells.² The kidney and skin also contain phytosphingoglycolipids although in much lower concentrations than in the small intestine. Phytoceramides form part of the water barrier lipids of the skin. 2hydroxytetracosanoyl-phytoceramide has recently been shown to have immunostimulating effects in humans.³ Phytoceramides have lately been studied in regards to their role in the central nervous system and have been found to have important functions in neuroprotection.⁴

Selected References:

- 1. K. Ryu et al. "Anti-scratching Behavioral Effects of N-Stearoylphytosphingosine and 4-Hydroxysphinganine in Mice" Lipids, Vol. 45 pp. 615-618, 2010
- 2. F. Omae et al. "DES2 protein is responsible for phytoceramide biosynthesis in the mouse small intestine" Journal of Biochemistry, vol. 379 pp. 687-695,
- 3. M. Sekiya et al. "A phytoceramide analog stimulates the production of chemokines through CREB activation in human endothelial cells" Int Immunopharmacol., vol. 11 pp. 1497-503, 2011
- 4. J-C. Jung et al. "Phytoceramide Shows Neuroprotection and Ameliorates Scopolamine-Induced Memory Impairment" Molecules, vol. 16 pp. 9090-9100, 2011

This product is to be used for research only. It is not intended for drug or diagnostic use, human consumption or to be used in food or food additives. Matreya assumes no liability for any use of this product by the end user. We believe the information, offered in good faith, is accurate.